

令和6年産 黒大豆収量調査野帳(脱粒後)

調査地点	定点	調査年月日	令和7年1月16日			※水分を15%換算、10a当りで計算					
調査項目		単位	畑井	福住	八上	大野	町ノ田	上小野原	合計	平均	割合
粗子実重			112.7	189.1	146.2	84.1	253.2	155.6	941.0	156.8	100.0%
くず粒			31.3	27.3	67.7	47.7	77.7	64.9	316.6	52.8	33.7%
		病虫害粒	2.7	2.8	2.3	4.0	2.3	8.4	22.5	3.8	2.4%
		破砕粒(割れ豆)	4.0	5.1	4.5	8.2	11.7	12.0	45.5	7.6	4.8%
		腐敗粒	3.7	0.9	2.7	4.8	5.9	7.0	24.9	4.2	2.6%
		はく皮・裂皮粒・猫ヒゲ粒	19.2	16.5	57.7	30.2	56.4	36.6	216.6	36.1	23.0%
		その他くず粒	1.7	2.0	0.5	0.6	1.4	1.0	7.2	1.2	0.8%
規格外粒			49.9	89.5	45.0	26.0	109.4	61.4	381.3	63.5	40.5%
		しわ粒	9.4	40.2	2.4	1.5	59.7	8.0	121.2	20.2	12.9%
		障子粒	18.3	35.0	16.4	2.7	40.3	32.8	145.5	24.2	15.5%
		茶斑粒	6.2	4.3	13.8	11.7	4.6	9.5	50.2	8.4	5.3%
		変形粒(未熟粒・扁平粒)	16.0	10.0	12.3	10.1	4.8	11.1	64.4	10.7	6.8%
		その他規格外粒	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0%
水分(3回測定)			16.7	16.2	19.7	24.3	15.6	18.4	110.9	18.5	
百粒重(2回測定)			65.6	57.4	66.7	73.1	54.6	63.9	381.3	63.6	
3L(11mm以上)			3.2	5.4	9.7	3.2	2.5	4.2	28.1	4.7	11.6%
2L(10mm以上)			20.1	42.2	17.6	5.9	33.0	17.8	136.5	22.7	56.1%
L(9.1mm以上)			7.5	22.1	5.9	1.0	26.3	6.6	69.4	11.6	28.6%
M(9.1mm未満)			0.7	2.7	0.2	0.2	4.4	0.7	9.0	1.5	3.7%
整粒重			31.5	72.4	33.4	10.3	66.1	29.3	243.1	40.5	25.8%