

令和7年度黒大豆定点調査成熟期生育とりまとめ(JA丹波ひかみ)

調査項目	単位	柏原	氷上	青垣	春日	山南	市島	定点平均
主茎長	センチ	73.7	71.0	58.8	78.5	75.0	74.7	71.9
全重	g/株	266.7	529.3	188.3	191.8	313.3	256.7	291.0
茎径	ミリ	16.6	20.6	15.4	19.1	18.2	16.8	17.8
主茎節数	節	21.0	22.0	15.5	17.2	17.7	17.8	18.5
一次分枝数	本/株	9.5	12.0	7.8	11.2	10.0	8.7	9.9
一粒莢数	莢/株	53.5	78.3	33.3	70.2	20.2	3.5	43.2
一粒莢重	g/株	43.3	73.5	22.3	56.7	16.2	2.5	35.7
二粒莢数	莢/株	69.0	62.0	34.0	78.5	10.2	0.8	42.4
二粒莢重	g/株	86.7	97.2	43.3	105.0	13.7	0.7	57.8
三粒莢数	莢/株	3.0	2.2	2.5	3.8	1.7	0.0	2.2
三粒莢重	g/株	5.0	4.8	4.5	6.7	2.8	0.0	4.0
外莢数	莢/株	14.5	20.3	7.0	12.7	9.3	15.8	13.3
外莢重	g/株	3.3	5.3	1.2	3.3	3.2	4.2	3.4
稔実莢数	莢/株	125.5	142.5	69.8	152.5	32.0	4.3	87.8
稔実莢重	g/株	135.0	175.5	70.1	168.3	32.7	3.2	97.5
総莢数	莢/株	140.0	162.8	76.8	165.2	41.3	20.2	101.1
総莢重	g/株	138.3	180.8	71.3	171.7	35.8	7.3	100.9

令和7年度黒大豆定点調査成熟期収量品質とりまとめ(JA丹波ひかみ)

調査項目	単位	柏原	氷上	青垣	山南	春日	市島	定点平均
地上部重	kg/10a	396.8	784.2	620.0	433.2	539.8	425.8	533.3
粗子実重	kg/10a	152.3	187.9	164.5	52.8	207.4	3.7	128.1
整粒重	kg/10a	89.3	71.3	74.5	15.0	93.4	0.8	57.4
規格外重	kg/10a	38.5	79.7	59.4	28.4	95.0	2.1	50.5
屑重	kg/10a	24.5	37.0	30.6	9.4	19.0	0.8	20.2
百粒重	グラム	59.7	67.3	57.5	56.3	59.6	48.1	58.1
3L率	%	1.3%	3.3%	4.5%	0.0%	2.7%	0.0%	2.0%
2L率	%	41.7%	68.7%	40.0%	25.3%	44.4%	20.0%	40.0%
L率	%	50.3%	24.9%	38.9%	59.7%	46.3%	60.0%	46.7%
M率	%	6.6%	3.1%	16.6%	15.1%	6.6%	20.0%	11.3%
整粒率	%	58.6%	37.9%	45.3%	28.5%	45.0%	21.7%	39.5%
規格外率	%	25.3%	42.4%	36.1%	53.7%	45.8%	56.5%	43.3%
しわ粒率	%	1.3%	2.9%	0.0%	0.6%	2.2%	0.0%	1.2%
障子粒率	%	16.6%	29.3%	18.8%	29.9%	12.9%	13.0%	20.1%
茶斑粒率	%	2.7%	5.3%	5.3%	5.0%	16.7%	4.3%	6.6%
変形粒率	%	4.7%	4.9%	12.1%	18.1%	13.9%	39.1%	15.5%
屑率	%	16.1%	19.7%	18.6%	17.9%	9.2%	21.7%	17.2%
病虫害粒率	%	0.8%	0.6%	2.5%	1.8%	0.3%	0.0%	1.0%
破碎粒率	%	0.3%	1.3%	2.2%	0.7%	1.2%	4.3%	1.7%
腐敗粒率	%	1.5%	3.4%	4.2%	0.3%	0.5%	4.3%	2.4%
裂皮粒率	%	13.5%	14.3%	9.7%	15.0%	7.2%	13.0%	12.1%

(粗子実重、整粒重、規格外重、屑重、百粒重については、水分15%換算値)