

班	地域	25株発生株数調査								本田 10回(5回往復)振りすくい取り調査										20回振	10株払い落とし調査											
		発生株数								頭数											頭数											
		いもち病	穂いもち病	紋枯病	ニカメイチュウ	イネミズゾウムシ	イネツトムシ	コブノメイガ	フタオビコヤガ	ヒメビウンカ	セジロウンカ	トビイロウンカ	ツマグロヨコバイ	フタオビコヤガ	カメムシ	ヒメビウンカ	セジロウンカ	トビイロウンカ	ツマグロヨコバイ		フタオビコヤガ	カメムシ										
									幼	成	幼	成	幼	成	幼	成	幼	成	カメムシ	幼	成	幼	成	幼	成	幼	成	幼	成	カメムシ		
1	下司											4				2			1													
	上河合	1									3					2			1	6	4											
	尾崎園出										4					2				40	15	45	15			3	1					
	生田								13	5	16	6			14	17			5	20	5	60	10			10	3					
2	宮前									3															1							
	小田										40	4						1			1	1			3							
	中持																			1	2				5							
	楠本									1		4			2							4										
3	木曽上									2		3						2	3		26	1										
	広石中									1		9						1														
	都志											2								1												
	下堺										2	1						27	1		5											
4	大野										3	3						3				1										
	由良										6	2									1											
	加茂	1								2	10	10									2											
	中川原	9							2		1	1											1									
平均株数		0.7	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.9	4.9	3.5	0.0	0.0	1.0	1.4			2.6	4.4	1.6	9.1	1.8	0.0	0.0	1.4	0.3					
発生地点率		18.8	0.0	0.0	0.0	0.0	0.0	0.0	12.5	37.5	43.8	87.5	0.0	0.0	12.5	25.0			50.0	31.3	31.3	56.3	37.5	0.0	0.0	31.3	12.5					
令和5年同時期調査結果		2.1	0.0	0.3	0.0	0.0	0.1	0.1	0.0	0.1	0.7	0.4	0.3	0.0	0.0	0.0	0.1			0.0	0.8	0.5	0.8	0.6	0.0	0.0	1.3	0.4				
		25.0	0.0	6.3	0.0	0.0	6.3	6.3	0.0	6.3	18.8	6.3	12.5	0.0	0.0	0.0	12.5			0.0	31.3	31.3	31.3	37.5	0.0	0.0	31.3	18.8				
令和4年同時期調査結果																																
令和3年同時期調査結果		4.8			0.0	0.0	0.0	0.0	0.0	0.6	1.9	0.9	0.5	0.0	0.1	0.3	0.4	0.0	0.1	0.3	2.9	1.4	3.0	0.4	0.0	0.0	1.8	0.7				
令和2年同時期調査結果		50.0			0.0	0.0	0.0	0.0	0.0	18.8	37.5	31.3	25.0	0.0	6.3	25.0	25.0	0.0	12.5	18.8	81.3	31.3	56.3	25.0	0.0	0.0	62.5	18.8				
		0.0			0.3	0.1	0.0	0.0	0.0	0.3	0.4	0.8	0.9	0.0	0.0	0.6	0.8			8.1	1.5	0.1	1.6	0.6	0.0	0.0	2.3	0.1				
		0.0			12.5	6.3	0.0	0.0	0.0	6.3	18.8	18.8	31.3	0.0	0.0	12.5	18.8			62.5	31.3	12.5	50.0	37.5	0.0	0.0	50.0	6.3				

◎結果と考察

◎対 策