

班	地域	25株発生株数調査								本田 10回(5回往復)振りすくい取り調査										20回振	10株払い落とし調査												
		発生株数								頭数											カメムシ	頭数											
		いもち病	穂いもち病	紋枯病	ニカメイチュウ	イネミズゾウムシ	イネツトムシ	コブノメイガ	フタオビコヤガ	ヒメトビウンカ	セジロウンカ	トビイロウンカ	ツマグロヨコバイ	フタオビコヤガ	ヒメトビウンカ	セジロウンカ	トビイロウンカ	ツマグロヨコバイ	フタオビコヤガ			カメムシ											
									幼	成	幼	成	幼	成	幼	成	幼	成		幼	成	幼	成	幼	成	幼	成	幼	成				
1	下司																																
	上河合	1																				4											
	尾崎園出																			1			1										
	生田															1					1	1					1						
2	宮前									2											2		3										
	小田																				17	8	1				2						
	中持										1	1									4	6	1										
	楠本																		18			2	2				1						
3	木曽上											2											4										
	広石中																																
	都志											1											2										
	下堺																		6			2											
4	大野																																
	由良											1																					
	加茂									6	5		2							2													
	中川原										1																						
平均株数		0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	0.1	0.4	0.0	0.0	0.0	0.1			1.6	1.5	0.9	1.3	0.2	0.0	0.0	0.2	0.1					
発生地点率		6.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6.3	18.8	6.3	31.3	0.0	0.0	0.0	6.3			18.8	25.0	18.8	56.3	12.5	0.0	0.0	12.5	6.3					
令和6年同時期調査結果		0.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.9	4.9	3.5	0.0	0.0	1.0	1.4			2.6	4.4	1.6	9.1	1.8	0.0	0.0	1.4	0.3					
		18.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	12.5	37.5	43.8	87.5	0.0	0.0	12.5	25.0			50.0	31.3	31.3	56.3	37.5	0.0	0.0	31.3	12.5					
令和5年同時期調査結果		2.1	0.0	0.3	0.0	0.0	0.1	0.1	0.0	0.1	0.7	0.4	0.3	0.0	0.0	0.0	0.1			0.0	0.8	0.5	0.8	0.6	0.0	0.0	1.3	0.4					
		25.0	0.0	6.3	0.0	0.0	6.3	6.3	0.0	6.3	18.8	6.3	12.5	0.0	0.0	0.0	12.5			0.0	31.3	31.3	31.3	37.5	0.0	0.0	31.3	18.8					
令和4年同時期調査結果																																	
令和3年同時期調査結果		4.8			0.0	0.0	0.0	0.0	0.0	0.6	1.9	0.9	0.5	0.0	0.1	0.3	0.4	0.0	0.1	0.3	2.9	1.4	3.0	0.4	0.0	0.0	1.8	0.7					
		50.0			0.0	0.0	0.0	0.0	0.0	18.8	37.5	31.3	25.0	0.0	6.3	25.0	25.0	0.0	12.5	18.8	81.3	31.3	56.3	25.0	0.0	0.0	62.5	18.8					

- ◎結果と考察

昨年と比べて、病虫害の発生率は少ない。
一部の地域で、畦畔すくい取りによる斑点米カメムシを確認。(楠本では、イネカメムシの幼虫も確認)
- ◎対 策

畦畔の草刈りを行い、出穂前にカメムシ防除を行う。
- ◎今後の管理

圃場内で出穂が7割を超えているところは、品質低下を防ぐために夕方に入水し、間断かん水を行ってください。