

班	地域	25株発生株数調査								本田 10回(5回往復)振りすくい取り調査										20回振 カメムシ	10株払い落とし調査																
		発生株数								頭数											頭数																
		いもち 病	穂い もち病	紋枯 病	ーカメ イチュ ウ	イネミ ズソウ ムシ	イネツ トムシ	コブノ メイガ	フタオ ビコヤ ガ	ヒメトビウンカ	セジロウンカ	トビイロウンカ	ツマグロヨコバイ	フタオビコヤガ	ヒメトビウンカ	セジロウンカ	トビイロウンカ	ツマグロヨコバイ	フタオビコヤガ																		
										幼	成	幼	成	幼	成	幼	成	幼	成		幼	成	幼	成	幼	成	幼	成	カメムシ								
1	下司																																				
	北山									3			1																								
	尾崎園出生田				3						1																										
2	宮前									16	1		13																								
	小田	2								1	2	1	1																								
	中持									10	3																										
	楠本									6	3																										
3	木曽上	2											1																								
	広石中											3																									
	都志											2	2																								
	下塚									6	2																										
4	新村																																				
	由良																	1																			
	下内善											99							1																		
	中川原											70							1																		
平均株数		0.3		0.0	0.2	0.0	0.0	0.0	0.0	2.8	0.8	11.0	2.5	0.0	0.0	0.0	0.2	0.0	0.0	0.0	1.1	0.3	10.8	0.1	0.0	0.0	0.3	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
発生地点率		12.5		0.0	6.3	0.0	0.0	0.0	0.0	43.8	37.5	37.5	43.8	0.0	0.0	0.0	18.8	0.0	0.0	0.3	18.8	18.8	56.3	6.3	0.0	0.0	25.0	6.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
令和7年同時期調査結果		0.0		0.1	0.0	0.0	0.0	0.0	0.0	0.4	1.3	0.1	0.1	0.0	0.0	0.0	0.1	0.0	0.0	0.7	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
		0.0		6.3	0.0	0.0	0.0	0.0	0.0	31.3	56.3	6.3	6.3	0.0	0.0	0.0	6.3	0.0	0.0	18.8	31.3	6.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
令和6年同時期調査結果		0.4		0.0	0.0	0.0	0.0	0.0	0.0	3.0	1.9	2.8	1.8	0.0	0.0	0.4	0.3	0.0	0.4	2.6	1.3	2.8	0.8	0.0	0.0	0.4	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0			
		12.5		0.0	0.0	0.0	0.0	0.0	0.0	50.0	50.0	43.8	62.5	0.0	0.0	12.5	31.3	0.0	0.0	12.5	50.0	31.3	25.0	43.8	0.0	0.0	18.8	12.5	0.0	0.0	0.0	0.0	0.0	0.0			
令和5年同時期調査結果		1.7		0.0	0.0	0.0	0.0	0.1	0.0	0.6	0.2	0.8	0.3	0.0	0.0	0.1	0.6	0.0	0.0	1.1	0.3	3.6	0.4	0.0	0.0	1.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0			
		18.8		0.0	0.0	0.0	0.0	6.3	0.0	25.0	18.8	25.0	31.3	0.0	0.0	12.5	25.0	0.0	0.0	25.0	25.0	31.3	12.5	0.0	0.0	37.5	18.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0			
令和4年同時期調査結果		1.1		1.8	0.0	0.0	0.0	0.0	0.0	3.9	1.5	18.5	7.3	0.0	0.0	3.2	5.7	0.0	0.0	1.1	2.9	0.5	3.1	1.6	0.0	0.0	0.8	0.1	0.0	0.0	0.0	0.0	0.1				
		18.8		43.8	0.0	0.0	0.0	0.0	0.0	18.8	25.0	81.3	81.3	0.0	0.0	18.8	31.3	0.0	0.0	62.5	56.3	25.0	56.3	50.0	0.0	0.0	37.5	6.3	0.0	0.0	0.0	6.3					
令和3年同時期調査結果		0.8		1.4	0.0	0.0	0.0	0.0	0.0	3.6	0.1	3.8	0.3	0.0	0.0	1.9	0.6	0.0	0.0	0.2	1.4	0.3	7.6	3.1	0.0	0.0	0.7	0.8	0.0	0.0	0.0	0.0	0.0				
		37.5		18.8	0.0	0.0	0.0	0.0	0.0	25.0	12.5	31.3	18.8	0.0	0.0	43.8	25.0	0.0	0.0	18.8	37.5	12.5	62.5	31.3	0.0	0.0	25.0	37.5	0.0	0.0	0.0	0.0	0.0				

◎結果と考察

- 1
- 病害に関しては一部いもち病やゴマ葉枯病が確認された。また、紋枯れの発病も確認された。
- 2
- 害虫では、全体的にウンカの発生が確認された。

◎対 策

- 1
- 病害虫防除においては出穂前防除を必ず行う。
- 2
- 出穂2週間前の畦畔の草刈りによりカメムシが本田への飛び込みを抑える。